



Low biomass production limits cover crop effects on soils

March 30, 2020

Cover crops at termination time in spring at Mead, NE. Photo courtesy of Sabrina Ruis.

Cover crops at termination time in spring at Mead, NE. Photo courtesy of Sabrina Ruis.

Cover crop impacts on soil properties depends on cover crop productivity. Planting cover crops early and in a diverse mix of species could be an option to boost biomass production and enhance benefits to soils. However, the impacts of early planting and species mixes on soil properties are not well understood.

A new article in *Agronomy Journal* investigates how broadcasting cover crops pre-harvest or drilling post-harvest affected biomass production and soil properties after four years. Cover crops were cereal rye, a mix of rye, legumes, and brassicas, and a no-cover-crop control. These were studied under no-till rainfed and irrigated continuous corn and corn-soybean rotations at three sites in the eastern Great Plains.

Cover crop biomass was low ($<1 \text{ Mg ha}^{-1}$) but was usually greater under pre-harvest than post-harvest establishment. Rye and mixed cover crops produced similar amounts of biomass. The low amounts of produced cover crop biomass led to few changes in soil properties that varied by site and specific soil property. Overall, early planted cover crops and cover crop mixes had minimal effects on soil properties in this region after four years, indicating the need for further research on cover crop impacts.

Dig Deeper

Ruis, S.J., Blanco-Canqui, H., Elmore, R.W., Koehler-Cole, K., Ferguson, R.B., Francis, C.A., & Shapiro, C.A. (2020). Impacts of cover crop planting dates on soils after four years. *Agronomy Journal*, 112. <https://doi.org/10.1002/agj2.20143>

[More science articles](#)

[Back to issue](#)

[Back to home](#)

Text © . The authors. CC BY-NC-ND 4.0. Except where otherwise noted, images are subject to copyright. Any reuse without express permission from the copyright owner is prohibited.